

Surface gravity and Hawking temperature from entropic force viewpoint

Ee Chang-Young,^{*} Myungseok Eune,[†] and Kyoungtae Kimm[‡]

*Department of Physics and Institute of Fundamental Physics,
Sejong University, Seoul 143-747, Korea*

Daeho Lee[§]

Basic Science Research Institute, Sogang University, Seoul 121-742, Korea

Abstract

We consider a freely falling holographic screen for the Schwarzschild and Reissner-Nordström black holes and evaluate the entropic force à la Verlinde. When the screen crosses the event horizon, the temperature of the screen agrees to the Hawking temperature and the entropic force gives rise to the surface gravity for both of the black holes.

Keywords: entropic force, Hawking temperature, surface gravity

^{*}cylee@sejong.ac.kr

[†]younms@sejong.ac.kr

[‡]helloktk@naver.com

[§]dhleep@sogang.ac.kr

Hawking radiation results from the quantum effect of fields in a classical geometry with an event horizon [1]. Thus the study of Hawking radiation may provide a clue in constructing the theory of quantum gravity. Considering its importance, it may be useful to have various interpretations of the effect from different angles, which may lead to a further understanding of the nature of black holes.

There have been several ways of providing the interpretation of Hawking radiation. One of them is the proposal suggested by Robinson and Wilczek [2], which is that Hawking radiation plays the role of preserving general covariance at the quantum level by canceling the diffeomorphism anomaly at the event horizon (see also [3, 4]). The flux of Hawking radiation can be also obtained through the scattering analysis and there have been the studies of the grey body factor for various black holes to calculate the Hawking temperature [5–14].

Recently, it has been suggested by Verlinde that gravitational interaction can be interpreted as a kind of entropic force through the holographic principle and the equipartition rule [15]. This entropic formulation of inertia and gravity has been used to study thermodynamics at the apparent horizon of the Friedmann-Robertson-Walker universe [16], Friedmann equations [17], Newtonian gravity in loop quantum gravity [18], holographic dark energy [19–21], an extension to Coulomb force [22], entropic corrections to Newton’s law [23], the Planck scale effect [24], and the gauge/gravity duality [25]. There have been many works for the entropic force in the cosmological models [26–31] and the black hole backgrounds [32–37]. In this paper, we study the Hawking temperature and surface gravity by calculating the entropic force on the holographic screen for spherically symmetric and static black holes.

We consider a test particle with mass m which approaches the holographic screen from a distance Δx (see FIG. 1). In this case the change of entropy associated with the information on the boundary is assumed to be [15]

$$\Delta S = 2\pi m \Delta x, \quad (1)$$

where we have taken the natural units of $c = \hbar = k_B = 1$. From the thermodynamic laws, the entropic force F acting on the test particle is given by [15, 38]

$$F \Delta x = T \Delta S, \quad (2)$$

where T is the temperature on the screen. Substituting Eq. (1) into Eq. (2), the entropic




FIG. 1: A test particle with mass m approaches the holographic screen, which is the boundary ∂V of the sphere with volume V with radius r . Δx is the distance that the particle moves.

force can be written as

$$F = 2\pi mT. \quad (3)$$

It is assumed that the total number N of the fundamental bits is proportional to the area A of the screen, *i.e.*,

$$N = \frac{A}{G}, \quad (4)$$

where G is a given constant for the moment and will be identified as the Newton's constant later on. When a total energy E is divided uniformly over the bits N , the temperature is determined by the equipartition rule [15]

$$E = \frac{1}{2}NT, \quad (5)$$

as the average energy per bit.

Now, we consider the Schwarzschild-like black holes which include both Schwarzschild and Reissner-Nordström (RN) black holes with the line element given by

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2d\Omega_2^2, \quad (6)$$

where $f(r)$ vanishes at the event horizon r_H .

In order to consider a freely falling holographic screen, we introduce the Painlevé-Gullstrand (PG) line element [39] of the Schwarzschild-like geometry (6). The ingoing (+) and outgoing (−) PG coordinates can be written as

$$ds^2 = -d\tau^2 + \left(dr \pm \sqrt{1-f}d\tau\right)^2 + r^2d\Omega_2^2, \quad (7)$$

or, equivalently,

$$ds^2 = -f d\tau^2 \pm 2\sqrt{1-f} d\tau dr + dr^2 + r^2 d\Omega_2^2. \quad (8)$$

Note that constant-time slices are completely flat in this coordinates system. The PG coordinates are related to the Schwarzschild-like coordinates by

$$d\tau = dt \pm \frac{\sqrt{1-f}}{f} dr. \quad (9)$$

The conserved energy of the gravitational field over a spacelike hypersurface V at a certain time is given by the Komar integral [40]

$$E(V) = \frac{1}{4\pi G} \oint_{\partial V} \nabla^\mu \xi^\nu n_\mu \sigma_\nu dA, \quad (10)$$

where n_μ and σ_μ are the unit normal vectors perpendicular to the hypersurface V and to its boundary ∂V with constant radius r , respectively. The normal vectors satisfy $n_\mu n^\mu = -1$ and $\sigma_\mu \sigma^\mu = 1$. And ξ^α is a timelike Killing vector field which satisfies the Killing equation of $\nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu = 0$.

For $r > r_H$, the Killing vector and the unit normal vectors are given by $\xi^\mu = \delta^\mu_\tau$, $n_\mu = -\delta^\tau_\mu$, and $\sigma^\mu = \delta^\mu_r$ in the PG coordinates. Then, from Eq. (10), the energy is given by

$$E = \frac{r^2}{2G} f'(r), \quad (11)$$

where $'$ denotes a differentiation with respect to r . The entropy defined on the freely falling holographic screen located outside the horizon is given by

$$S = \frac{A}{4G}. \quad (12)$$

The above relation can be restated as

$$N = 4S \quad (13)$$

via Eq. (4). Using the equipartition rule for the energy E in Eq. (5) and the relation (13), we get the following relation [41, 42]

$$T = \frac{E}{2S}. \quad (14)$$

Then, from Eqs. (11) and (12) the temperature on the holographic screen with the constant radius r is given by

$$T(r) = \frac{1}{4\pi} f'(r). \quad (15)$$

At the horizon, the surface gravity κ is defined with the Killing vector ξ^μ by the following relation

$$\xi^\mu \nabla_\mu \xi^\alpha = \kappa \xi^\alpha. \quad (16)$$

In the Schwarzschild-like geometry, $\kappa = \frac{1}{2}f'(r_H)$. Thus, the temperature (15) evaluated at the horizon ($r = r_H$) is related to the surface gravity as follows:

$$T(r_H) = \frac{1}{4\pi}f'(r_H) = \frac{\kappa}{2\pi}. \quad (17)$$

Since the Hawking temperature T_H is related to the surface gravity by $T_H = \frac{\kappa}{2\pi}$ [43], the temperature at the horizon $T(r_H)$ is the same as the Hawking temperature.

Finally, the entropic force is obtained from Eqs. (3) and (15) as

$$F = 2\pi m T(r) = \frac{1}{2}m f'(r), \quad (18)$$

where m is the mass of the test particle. For the Schwarzschild case $f(r) = 1 - 2GM/r$, the entropic force becomes the Newton force

$$F = \frac{GmM}{r^2}, \quad (19)$$

where M is the mass of the black hole and G is identified as the Newton's gravitational constant. For the RN case $f(r) = 1 - 2GM/r + GQ^2/r^2$, the entropic force becomes

$$F = \frac{GmM}{r^2} - \frac{GmQ^2}{r^3}, \quad (20)$$

where M and Q are the mass and charge of the black hole, respectively. Note that the force in Eq. (20) contains the gravitational effect only, since the test particle's charge does not play any role in this consideration. Via the metric, only the geometry which is determined by the energy of the system plays a role here. At the event horizon ($r = r_+$), the entropic force in the RN black hole becomes the usual expression

$$F = \frac{m(r_+ - r_-)}{2r_+^2} = m\kappa,$$

where $r_\pm \equiv GM \pm \sqrt{G^2M^2 - GQ^2}$.

In Ref. [22], the RN black hole was considered. There the motivation came from a mismatch between the Hawking temperature and the energy in the entropic force formulation. However, the energy used there contains only the gravitational mass not including the electromagnetic energy. On the other hand, we obtain the energy using the Komar integral with the RN metric, which contains both mass and charge contributions to the energy. This yields the correct relation in our case.

Acknowledgments

We thank Youngone Lee for helpful discussions. This work was supported by the National Research Foundation (NRF) of Korea grants funded by the Korean government (MEST) [R01-2008-000-21026-0 and NRF-2009-0075129 (E. C.-Y. and K. K.), NRF-2009-351-C00109 (M. E.), and NRF-2009-351-C00111 (D. L.)].

-
- [1] S. W. Hawking, Commun. Math. Phys. **43** (1975) 199 [Erratum-ibid. **46** (1976) 206].
 - [2] S. P. Robinson and F. Wilczek, Phys. Rev. Lett. **95** (2005) 011303 [arXiv:gr-qc/0502074].
 - [3] R. Banerjee and S. Kulkarni, Phys. Lett. B **659** (2008) 827 [arXiv:0709.3916 [hep-th]]; Phys. Rev. D **79** (2009) 084035 [arXiv:0810.5683 [hep-th]].
 - [4] R. Banerjee, Int. J. Mod. Phys. D **17** (2009) 2539 [arXiv:0807.4637 [hep-th]].
 - [5] K. Ghoroku and A. L. Larsen, Phys. Lett. B **328** (1994) 28 [arXiv:hep-th/9403008].
 - [6] S. S. Gubser and I. R. Klebanov, Phys. Rev. Lett. **77** (1996) 4491 [arXiv:hep-th/9609076].
 - [7] J. M. Maldacena and A. Strominger, Phys. Rev. D **55** (1997) 861 [arXiv:hep-th/9609026].
 - [8] I. R. Klebanov and S. D. Mathur, Nucl. Phys. B **500** (1997) 115 [arXiv:hep-th/9701187].
 - [9] M. Cvetič and F. Larsen, Nucl. Phys. B **506** (1997) 107 [arXiv:hep-th/9706071].
 - [10] H. W. Lee and Y. S. Myung, Phys. Rev. D **58** (1998) 104013 [arXiv:hep-th/9804095].
 - [11] W. T. Kim and J. J. Oh, Phys. Lett. B **461** (1999) 189 [arXiv:hep-th/9905007].
 - [12] W. T. Kim, J. J. Oh and K. H. Yee, Phys. Rev. D **66** (2002) 044017 [arXiv:hep-th/0201117].
 - [13] G. Clement, J. C. Fabris and G. T. Marques, Phys. Lett. B **651** (2007) 54 [arXiv:0704.0399 [gr-qc]].
 - [14] W. Kim and J. J. Oh, J. Korean Phys. Soc. **52** (2008) 986 [arXiv:0709.1754 [hep-th]].
 - [15] E. P. Verlinde, arXiv:1001.0785 [hep-th].
 - [16] F. W. Shu and Y. Gong, arXiv:1001.3237 [gr-qc].
 - [17] R. G. Cai, L. M. Cao and N. Ohta, arXiv:1001.3470 [hep-th].
 - [18] L. Smolin, arXiv:1001.3668 [gr-qc].
 - [19] M. Li and Y. Wang, arXiv:1001.4466 [hep-th].
 - [20] D. A. Easson, P. H. Frampton and G. F. Smoot, arXiv:1002.4278 [hep-th].
 - [21] U. H. Danielsson, arXiv:1003.0668 [hep-th].

- [22] T. Wang, arXiv:1001.4965 [hep-th].
- [23] L. Modesto and A. Randono, arXiv:1003.1998 [hep-th].
- [24] S. Ghosh, arXiv:1003.0285 [hep-th].
- [25] Y. Zhao, arXiv:1002.4039 [hep-th].
- [26] C. Gao, arXiv:1001.4585 [hep-th].
- [27] Y. Zhang, Y. g. Gong and Z. H. Zhu, arXiv:1001.4677 [hep-th].
- [28] Y. Wang, arXiv:1001.4786 [hep-th].
- [29] S. W. Wei, Y. X. Liu and Y. Q. Wang, arXiv:1001.5238 [hep-th].
- [30] Y. Ling and J. P. Wu, arXiv:1001.5324 [hep-th].
- [31] D. A. Easson, P. H. Frampton and G. F. Smoot, arXiv:1003.1528 [hep-th].
- [32] Y. S. Myung, arXiv:1002.0871 [hep-th].
- [33] Y. S. Myung and Y. W. Kim, arXiv:1002.2292 [hep-th].
- [34] Y. X. Liu, Y. Q. Wang and S. W. Wei, arXiv:1002.1062 [hep-th].
- [35] R. G. Cai, L. M. Cao and N. Ohta, arXiv:1002.1136 [hep-th].
- [36] Y. Tian and X. Wu, arXiv:1002.1275 [hep-th].
- [37] X. Kuang, Y. Ling and H. Zhang, arXiv:1003.0195 [gr-qc].
- [38] S. Hossenfelder, arXiv:1003.1015 [gr-qc].
- [39] M. Visser, Class. Quant. Grav. **15** (1998) 1767 [arXiv:gr-qc/9712010].
- [40] S. M. Carroll, *Spacetime and geometry: An introduction to general relativity* (Addison-Wesley, San Francisco, 2004).
- [41] T. Padmanabhan, Class. Quant. Grav. **21** (2004) 4485 [arXiv:gr-qc/0308070].
- [42] T. Padmanabhan, arXiv:0911.5004 [gr-qc]; arXiv:0912.3165 [gr-qc].
- [43] G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15** (1977) 2752.
- [44] F. Caravelli and L. Modesto, arXiv:1001.4364 [gr-qc].